# The expert Gardener:

OR,

A Treatise containing certaine necessary, secret, and ordinary knowledge, in Grafting and Gardening; with divers proper new Plots for the Garden.

Also sundry expert directions to know the time and season when to sow and replant all manner of Seeds. With divers remedies to destroy Snailes, Canker-wormes, Moths, Garden-Fleas, Earth-wormes, Moles, and other Vermine.

Faithfully collected out of sundry Dutch and French Authors.



LONDON, Printed by William Hum. 1654. The expers Gradence:

Wilder of the state of the stat

The state of the s



LOMDON, Priored by William. 165%



#### Certaine common Instructions bow the Stumpe must be chosen whereupon you will graffe or plant.



Very diligent Housholder who will plant, should use thereto a convenient place, to the end, that the wild beast chaw not, nor paire the plants; or if they be young, wholly eat in pieces; which to avoid, is needfull to

be in a towne or closed Orchard, where there is not too much shadow, but a sweet ground well muckt, rilled and turned.

Every Plant will have foure things.

First moistnesse, so that the seeds or stumpe bee

moist or green.

Secondly, a convenient place, which hath such earth as will lightly be rubbed to pouder, and that Sun may come to it; for where there is filthy lome, a lean ground, or sandy, dry, burnt, or salt ground, there is nothing good to be planted, to have any continuance; neverthelesse where the ground is lean, there you must give more dung; in a fat ground not so much. Take heed the ground be not too moist.

#### The expert Gardener.

nor too dry ; and muck the trees with hogs dung. Thirdly, a mediate water or neurithing moisiness. therefore be those Orchards best which are scituated between two waters; for those that are placed by a water fide, remaine feill young and fruitfull, and have commonly the back smoother and thinner than the others. And those trees are more fruitful then others which are planted in a vally, or in the lower part of a deep hill; for from those hils may come to them nourishment and moinness, and the ground which is fo scituated is very fruitfull. But he that cannot get for his trees fuch a ground, must with all diligence feek to bring to his trees a little spring or pond, of which the trees may fometimes find fome reviving. and if you may not have any of those; and have a garden who by it felfe is naught, the trees wil grow with thick roots, which hindereth the growing of them,

Fourthly, the air is required, which must be agreeable to them, and of complexion to bear; for there be some trees that doe prosper in all aires, to wit, apple, and peare, cherry and plum-trees. Some will have a cold air, to wit, chesnut-trees; and some a very warmaire, as the palme and pepper trees: therefore they be rare with us. That plant which hash these sour things shall prosper; and if they want one or more of shese four things, they will decay and their prosper.

ring periff.

and drieth them at length.

At what time trees ought to be planted and fer.

A LL kind of trees may be planted, transported, and cut in March, but it is better to turn them in Odober, for them the frost hurtesh them not for much

much as at other times; for learned men fay, that in dry Towns and warm Countries they plant in October or November, and that in moift Towns and cold vallies, they plant in February or March: in none other time may you plant or graffe. When you will plant or fet agains wild flumps, if there be any thing broken at the root, cut it off. Every plant must be fet two foot one from another, or at the least one foor, especially when they should beare strong fruites: likewise when thou wilt set strong seeds, as nuts, almonds and peaches. When a man will plant two stumps, so must they be of two yeare old, except the vine.

These things you must understand of those plants or stumps which are planted with roots.

## How the sumps and plants must be prepared and dressed, which you will plant.

The plant or sprout you must cut round about, so that you leave the very end of it, and put it then into a hole, but if the stump be great, cut it clean off, and then put only the undermost part into a hole, long or shore as you will; but if you find two stumps grown together, you may cut the lesser away. And above all things, you must take heed that the sprout grow upright, and if it will not, you must constraine it, and tie it to a stick.

Hero follow certaine instructions how the trees must be kept, and how you must tabour them.

Some trees will have a fat ground, as Figge trees and Mulberrietrees, and some leane ground, but

A 3

all trees be in that point equall, that they will have in the top dry ground, and in the bottome moist earth.

2. In Harvest you must uncover the roots of the trees so deep, that they may partly be seen, and lay dung upon them, which dung must be dissolved of raine in the ground, that it may come to the roots, which mucking giveth good increase to the roots.

3. If the ground wherein the trees stand bee too fandy, then mix among it faire and new lome; and if it bee too lomy, then mix amongst it sand in place of mucke, the which you must not only doe hard by the tree, but also four or five foot off from it round about the tree, according as the tree is in bignesse, or that the roots are large and great.

Such diligence, giveth to the trees great help, for their nourishment and strength is thereby renewed. Hereaster you shall understand, whereby to know

the fruitfull soile.

4. In the fat ground, the stumps whereupon you wil graff, must be lest long, but in lean ground short.

5. The plants of trees from their youth, till three years must not be cut nor shred, but they may bee transported, and if they be too weak you may pricke slicks next unto them.

6. Diligent regard must be taken, that no sprouts spring out of the stump, which might take the nou-rishment from the tree sprouts, and those boughes which spring from the root of the tree at the first planting.

- 7. When thou perceivest the young trees to wax weake, then uncover the roots and put other fresh

ground to them.

8. If the ground be neither too fost nor too hard, then may you chuse all kind of stumps in February for to plant, when the green juice is dispersed in the bark; bur when the ground is too hard, then the sweet holes or pores of the root doe remaine closed and stopped, so that they cannot draw to them their nou-rishment, such hardness of the ground or earth hindereth the aire and moistnesse which commeth from beneath upward, for it cannot be pierced of the soft sprouts, with the small heat which is beneath, therefore you must come to help them with a spade, for with a plough you will never come to an end, because of the root.

g. There is great diligence to be taken for preferving of the trees, when they begin to grow great? to scrape from the barke all rudenesse which is done, when you take from them all superfluity and sprouts which come out of the tree. You may cut them in February:

10. It is good for the trees to muck them often,

and moderatively to water their roots.

Also to cleave the roots, and lay stones into them, to the end they may revive againe of the drinesse which they have suffered, or of the barrennesse of the ground, or when the young planted trees for the great heat will perish. Also when immoderate heat is, then you must help them with turning of the ground, and with watering, but the water wherewith you should water them, must not be altogether fresh nor cold, or newly, drawne one of the spring; but out of a ditch, pond, or well, of any other sould ditch water, or with spring water, which hath stood long in the Sunne, or put a little dung in the water, and stire

it once or twice well about, and the water will be fat wherwith water your trees. You may also keep them with shadowes and straw from the heat: or else put (in great heat) fat green herbs at the stump, tempred with loame: some anoint the stumpe (toward the South or Mid-day) with chalke, some with oile, or with any other ointment that cooleth.

11. When you would transpose a plant, or have wilde Rumps digged out to plant again, then marke the part which standard towards the South of Mid-

day, and put it so againe when you graffe it.

# How to keep plants, flumps, or trees, from the wild beaft, that they burt them not.

Where the path of the beaft is free and remedilesse, there must be put poles, and with thornes the same young trees must be inclosed.

#### That the Detres Spoile them not.

TAke the pille of a Deere and anoint the Tree

#### That the Hares due not burt them;

SPet in thy hand, and anoint the sprouts therewith, and no Hare will hurt them.

Here

#### The expert Gardener?

#### Here follow fome infruttions of graffing.



Irst, you must know that imping, graffing, and setting, is all one thing.

The imping fprouts must be young and new, with great bodies and many eyes: for

where many and great bads be, that is a token, that

is of a ftrong fruit.

2. The imping sprouts must be broken off at the Sun rising, although that those of the other side broken off grow likewise; yet those of the other side are most naturall and temperate of heat; Some country clownes believe, that if you in cutting the sprouts turne them upside downe, that they will never grow right, but be crooked.

3 All graffing and imping is done by putting one into

4. Over-young imps (which are so weak that they will breake before they be put into the earth, or into the stump) are haught, and therefore they may not

be imped or fer,

5. When you imperipon a houle of truit tree, the fruit will be far better: But if you cut of a Garden tree a branch, and imp it hat one of his own sprouts, it will bring forth fruit of another talle, forme, and bigreffe; for imping maketh all the diversities in peares, apples, and other fruits.

6. It is far better to impe low in the flump, than in the top in the high branches: yet neverthelesse if you will make of wild apple trees garden trees, you

may importeen upon the top.

7. In great trees which have a great bark, it is not fo good to impe, for they take not to them so easily the voins of the roots which grow out of the young sprouts, because of their hardnesse, and especially when the imping sprouts are too weak. Wherefore they which graffs trees must seek small and young stumps, wherein they since much liquor and little hardnesse, and which may endure the banding.

8. It is bert imping or graffing when the liquor is in the bark, Pyou have a great tree upon the which youwould impeland hath many branches, you may quettern alloff, land impelant the flumps all kind of bought, lach as you pleafe; but if the tree be overled, for has her boughts be rancked, and her most melle confumed, then cut the tree clean off, and let abo flumps frid as whole years a fluctuard rake the oun.

fprouts which are sprung out of that stumpe, and graffe them, and cast the others away. Such a stump is like to bear, and therfore nourish as many sprouts as you please: but if it be a wild stump, graft garden

fprouts upon it.

9. If you graffe a sprout or bough upon a Hawthorne tree, that same bough will grow great, and
the stump will remaine small, therefore he that will
impeupon such a tree, see he cut it off by the roor,
then will the imped sprout and the stumpe grow all
of one thicknesse: but you must have still regard that
you impe kinde upon kinde, as apples upon apples,
peares upon peares: for he that graffeth strange upon
strange, as peares upon apples, and apples on pearer,
and such like, although it be done often for pleasures
sake, yet will it not last; for the natural nourishment
is so, that it will hardly nourish a strange kinde of
fruit.

10. The tree which is graffed in February, in his

fruits grow no worms nor maggots.

and will not grow streight and levell, then you must constrain them perforce, that they may grow orderly. Furthermore you must have a care to keep the prospering sprouts well with sticks from the wind, if they stand any thing high, and especially when they have stood a year or two, and where they are pricked in the stump, it is most needfull, as shall after appeare. And because there be many and divers waies to graffe, and know how wild stumps and trees are to be made garden trees, we thought it good to set some of them here down.

#### Divers fostions and mates of graffing there be:

HE that will extraordinarily graffe all manner of

ker another, the better, it will prosper.

The first fort of graffing is, when the sprout is prickt between the bark and the wood of the stump; which must be done in May or Aprill, when the bark may easily be loosed from the tree, and is done after this fort.

First take a stump or tree, and cut him off with a tharp Saw, knife or fuch like inftrument, where he is smoothest and clearest, and full of juice, and polish the place with the bark of the fame tree which was cut off. Afterward ty the flump with a picce of bark, and then prick a hole between the barke of the tree with a prick of bone, elderwood or Iron, fo that it cleave not, and then put in the place of the prick the forout, which you must have broken off a plaine and even tree, of a good kind, and one year old, which you shall know by this severy branch hath rinckled knots fike the joint of a mans finger, cut it at one fide under the knot, fo that you touch not the heart of the tree, and at the other fide you must fortly loose the barke, that the fp. oat may joyne very close to the flump, then pull out the prick, and take the sprout, and turn the green bank to the bank of the flumpe, fo that it may fland ftraight.

The sprout may be source or five fingers, or eight

at the most high above the stump.

Of this fort of imping, you may fee two, three, or more, according to the bigneffe of the flump, or as

he

lie can beare, provided alwaies that they fland at the

least the length of a finger one from another.

Afterward tie it fast (with bark) together, and put over it good muck, and tie over it a cloath, that no raine or aire may come between it and hurt it. This fort of imping is commonly used in stumps, which are great and old trees, whose barke is thicke and strong, as pple trees, pear trees, cherrytrees, and willow trees, on which are imped oftentimes apples, also on figge trees, and chestnut trees.

Such grafting is also done in high stumps, and branches, which be great, but they must be well-kept

from the wind, that it doe not break them.

After this fort you may graft many forts and kinds of pears upon one tree, but if you bring peares upon apples, or apples upon pears flumps, it will not last long, as afore is said.

The first way of grafting prospereth best, and hath a good continuance, there be many other forts of

grafting, as followeth.

another way of grafting is, when the stumps are clowen, and the sprouts afterwards are put in, the which doe as followeth:

Ake a young tree which is fount of the bigness of a finger, and cut it smooth and even, and eleave it in the midst, then take the sprout which you will impe, and cut it three square, and at the one side leave the barke uncut, and then turne the same bark outward at the stump; and the trans that I have taught, that the winder nor raine burt him not.

Otherwife. 2 01 101

When the flump is uncovered and clean burniffit

2012

at the foft place, then tie him faft, that he cleave no further than to the length of your foront, which you must graffe upon him, and then leave the prick in it. then make your sprout pointed like a prick, so that the middle be not touched, then put it into the cleft, having cleanled the hole first with the point of knife, to that one bark may touch the other, and ourward one wood another, to the end, the mediture may have the more easier his course, then pull out the bricke, and that which remaines open and bare between the cleft and the sprout, that bind well every where with the bark of the tree, or with hard preffing with a little fund, or with dung of an Oxe, or with waxe, or with a linnen cloth washed in waxe. that no raine, winde or worms may hurt it, This helpeth much to keep the moistnesse in, which commeth from the root, that it cannot breake out, but nourisheth the better the new plant; but when the flumps are great, they be cleaved after two waies. The first is, that you cut or cleave the tree with a knife at one fide only, even to the heart, and that you graft into it but one sprout. The other is, that you cleave it all over, and that you prick or graft on every fide one sprout, or one alone, and leave the ther fide without.

When the flump is but a little bigger, then the fprout must necessarily be cloven in two, and you must graft but one sprout into it, as is said in the be-

ginning.

This cleaving may be done in February, March, and Aprill, then it is good to cut them before they be greene, for to keepe them the better under the ground, in cold or moist places.

The

The third way of grafting.

His fort of grafting is very fubtill, witty, and

I ready; and is done as followerb.

Go to a smooth apple or peare tree, in April, when the trees get liquor, and feeke a branch which hath green eyes, and fee that the fame be leffe than your little finger, and teare it from the tree, and where you fee that the green sprouts will come off, there out them off wholly, and cleanse the middle thereof that the little red at the wood may turne about, and draw it not off, untill you come unto another good peare or apple tree, and feek there another branch of the fame bigneffe that the other was, and cut it off, and take from it, likewise the red, as far as you will put them again, and look where the branches join, that they may well fit together upon the top, and tie the' fame place gently and well with a little barke, behinde and before, that the water may not burt them: in the first year it bringeth forth leaves and branches, in the second floures, which you may breake off, for the sprout is yet too tender, so that it may bear no fruit, and in the third yeare it bringeth floures and fruit, and by this meanes you may graft divers kinds of peares and apples upon one tree. I have likewife fer fuch sprouts upon wild stumps, and they have prospered.

The fourth way of grafting is,

ther tree, like as a plaister is tied to a mans body:
this fort of grafting, is called in Latine Emplastrum.
We read of such a fort of grafting which is called in
Latine Abducellum, and it is much like unto this fort,
wherefore we will only speake of it, being done after
this fort.

When

The expert Gardener.

When you fee upon a great fruitfull bough a bud which will prosper without doubt, and wouldft faine plant it upon another tree, take a sharp knife, and lift the bark up two fingers breadth, that the bud be not hurt, then goe to another tree, upon the which you will graft, and put into a convenient place, a like hole into the bark, and put the same bud with the bark into it, and tie it with dung (or with a clout that hath lien in a dunghill) over the cut, that it may be kept from the outward damage of weather, and for an especiall nourishment and keeping of the inner juice: then cut off the branches round about it, that the mother may the better nourish the new sonne; within twenty days after take away the band, so that you see that the strange bud hath prospered, and joyned himdelfe with the tree. This may be done in March, when the bark cometheafily from the tree. Also in April, May, and June, and yet thee prospereth both before and after a time, when you may conveniently finde fuch buds.

This fort of planting prospereth best in a willow tree or such like, which is pierced through, and is done after this fort.

The fifth way.

When you pierce a willow stick with a sharp piercer, see that between every hole, be left the space of one foot, and prick therein branches a little scraped, and put the stick into a ditch, so that the branches stand upright, one part of the stick remaining over the earth; and within a year after take it out of the ditch, and cur the stick as under, so find you the branches full of roots, and put every one into a hole in the ground, and tis fit the holes were stopped with lome, or with wax.

Some do take in March a fresh Beech tree, which is of a mans thickneffe, and pierce him overthwart with maine and great holes and fmall holes, till unto the lowermost barke, or quite through then take forouts or boughes, which be as big and small, that they may fit into the holes; and when you will put them into the Beech flump, you must scrape the uppermoft barke off, untill the green, and no further: then the bough must remain into the Beech, the sprouts must stand a foot, or somwhat lesse asunder. then keep your Beech stumps with the sprouts in a fresh ground, and skant a foot deep, you must first maime the sprouts, that they may not flourish; then the next March enfuing, dig it out with the sprouts, and cut it afunder with a faw, and every block which is cut off with its branch, you must fet in a fresh ground, and so they will bring forth the fruit the fame yeare.

The fixth Way.

This way teaches how to graffe, that they may bring forth fruit the first years, which do as follows.

Pare an old stumpe of what kinde soever it be, the uppermost bark, till to the lower green barke, a span long or somewhat lesse, which do in Harvest in the wane of the Moone, and anoint it with Oxe dung and earth, and tye it with bark, and after in March when trees are transposed from one place to another then cut the same branch from the tree, and put it into the ground, and it will bring fruit the same yeer. I have seen that one hath prickt sticks on Albahar eve, in the earth, and hath pulled them out again upon Christman eve, and put boughes in the holes, and they have prospered and come out.

The feventh.

Pierce the top of a stump, which is not over small, and draw a bark through it, and maime it with a knife, as far as it standeth on the top, & in eight days after, poure water upon it, that the top of the stump may close. This must be done in harvest, and in the March following cut it off from the tree, and bruise the top, and put it with the same earth in other ground.

The eighth Way.

Will you graffe a tree that the fruit be without stones. Take a sprout and graff it into a great stump, with the thicker and lower part of the sprout, then take the upper or thinner end of the sprout, and cut it also fit to be graffed, and turne it downeward, and graffe it into the said stump; and when the sprout of both sides prospereth, cut it in the midst a sunder, so that which is grown right upward with the tree, the fruit of it hath stones, but that which was the top of the sprout that groweth contrary, brings forh fruit without stones. And if so be the turned sprout prosper, you must break off the other, to the end, that the turned sprout doe not perish, which you may try after this sort; for oftentimes it commeth and prospereth, and many times it is perished and spoiled.

How Cherries are to be graffed, that they may

Ill you make that Cherries grow without stones? pare a little Cherry tree at one year old at the stump, and cleave it as funder from the rop to the root, which do in May, and make an Iron sit to draw the heart or maxow from both sides of the tree; then

then tye it fast together, and anoint it with Ox dung or lome, and within a yeare after, when it is growne and healed, goe to another little tree which is of the same kind, and which hath not yet brought fruit, and graffe that same on the little tree, so shall that same tree bring his fruit without stones.

PLant a Vine tree next unto a Cherry tree, and when it groweth high, then pierce a hole into the Cherry tree right above it, that the hole be no bigger than the Vine is thick, and pare the upper bark of the Vine branch till unto the green, so farre as it must go through the tree, and look well to it that the branch of the Vine be not bruised, and wel anointed. You must not suffer any sprouts to come out of the Vine from the ground up, but unto the tree only, that which commeth out of the other side, let that same grow and bring fruit. Then the next March sollowing if the Vine prosper and grow fast into the tree, then

How a grape of a Vine may be brought into a glaffe.

Ill you make that a grape grow into a narrow glaffe? take the glaffe before the grape cast her bloud, or while she is little, and put her into the glasse, and shee will ripen in the glasse.

cut the Vine from the tree off, and anoint the place

with diligence, and it will bring fruit.

To graffe Medlers on a Peare tree.

If you graffe the branch of a Medler upon a Peare tree, the Medlers will be sweet and durable, so that you may keep them longer than otherwise.

#### The empere Gardener.

IF you will graft upon a wild flump, put the fprouts in Pikes bloud, and then graft them, and the fruit will be red.

Osbermife.

Take an apple branch, and graft it upon an alder flump, and the apples will be red. Likewise if you graft them upon cherry trees.

Ofthe Quince tree.

He Quince tree commeth not of any grafting, but you must plucke him out by the roots, and plant him againe into a good ground or earth,

Otherwise.

The Quince tree requireth a dry and fweet ground,

and he prospereth therein.

How to make that Quinces become great.

TAke a branch of a Quince tree when it father cast his bloud where a Quince groweth at, and put it into a pot, and set it into the ground, and set the Quince grow in it, and it will be very great.

And if you will shew some cunning therewith, cause to be made a pot which hath a mans face in the bortome of it, or any other picture whatsoever, and when the quinces have blossomed, then bow the branch, and put the Quince into the pot, and she will grow very big, in the shape of a man, which may also be done in Pompons, Mellons, Cucumbers, and other earthly fruites.

The conclusion of graffing.

OV It of all the forewritten causes (gentle reader) is evidently showne, that although every planting or grasting be better from like to like, and from kinde.

#### The expert Gardener.

kinde to kinde, yet neverthelesse ir agreeth also wish contrary kinds, as now is said wherefore he that will exercise and use the same, and try divers kindes, he may see and make many wonders.

#### What joy and fruit commet b of trees.

The first fruit.

The first is, that you plant divers & many kinds: for every housholder who hath care to his nourishment, with all diligence causeth oftentimes, such trees to be brought from forrein Countries.

The fecond.

The fecond is, when the trees be planted and fet orderly and pleafantly, they give no small pleafure to a man, therefore every one should cur his trees orderly, and he that cannot, should procure other men to doe it, which know how to doe it.

The third is of well [melling and spiced fruit.

Cleave a tree afunder, or a branch of a fruitfull tree, to the heart or pith, and cut a piece out of it, and put therein poudred spices, or what spice sever you will, or what colour you will defire, and tye a barke hard about it, and anoint it with lome and Ox dung, and the fruit will gett both the savour and colour according to the spice you have put in it.

How fowre fruits be made [weet,

Which tree beareth fowre fruites, in the fame pierce a hole a foot or fomewhat lesse above the root, and fill that with honey, and stop the hole with a haw-thorne branch, and the fruit will bee fweet.

C.3

Hom

The expert Gardener.

How trees ought to be kept when they wax old.

Hen trees lose their strength and vertue for age, and the branches break off for the weight of the fruit, or when they wax barren for lack of moisture, that they beare not fruit every year, but scant every other or third yeare, you must cut some of his heavy branches, which he can little nourish, which is done to the end he might keep some moistnesse to himselfe for his nourishment, or else the moistnesse would goe all into his branches.

Whereby you may mark whether you must give them, or take away from them branches, according to their nourishment, and as the earth where shee standeth can abide, that is, you must leave them so much as will nourish them, and no more, which if you doe not, the trees will bring so little fruit, that

your labour will not be recompensed.

Which cutting of trees may be done from the beginning of November till to the end of March, in warme countries. But it is more naturall to be done from the time that the leaves fall, till the time that they begin to grow green againe, except where the frost is very great and sharp.

How trees must be kept from divers sicknesses, and first bow to keep them from the Canker.

Hen the Canker commeth in any tree, he becommeth barren and dry, for it mounteth from the stumps into the top, and when it taketh a peare or apple tree, the bark will be black and barren thereabouts, which must be cut off with a knife, to the fresh wood, and then the place must be anointed with Oxe dung, and tied with barke, so that neither wind nor rain may hurt it.

Against

Against worms which must be driven out of the tree.

IT happeneth oftentimes, that the superfluities of moistnesse in the trees breaketh out like as sometimes to a man or beast between the slesh and skin: and when that beginneth to rot, wormes grow out of it, which takes his strength away: wherefore mark,

When the barke of a tree at any time swels, cut it presently open that the poison may runne out, and if you find already wormes in it, draw them out

with a little Iron book.

How the wormes are to be killed, if they be already grown into the tree.

I f you will kill the worms which grow in the tree, take Pepper, Lawrell, and Incense, and mingle all well together with good wine, and pierce a hole into the tree downeward, to the pith or heart of the tree, and poure this mixture into it, and stop it with a hawthorne, and the worms will dye.

Otherwife.

Take afties or dust and mingle it with sallet oyle, anoint the trees therewith, and the worms will dye.

Otherwise.

Take powdered Incense when you graffe, and bring it between the barke of the Rump, which you will.

graffe, and no wormes will eat the fruit.

When a tree in many places becommeth changeable because of wormes, or superstions humours, Cleave the tree at some end from the top of the stump to the earth, that all the soule liquors may come out and dry. Also when a tree becomes sick because of evill humours or fault of ground, so that he becommeth worme-eaten, or brings no fruit, take the earth away from the root, and put other sweeter in the place, and pierce a great hole in the stump, and put therein a pin of Oake, and it helpeth.

A remedy against Caterpillers.

A LL kind of Caterpillers which eat the green, and blostomes of the tree, doe hurt them very

much, so that thereafter may come no fruit.

Therefore their eggs which lye hidden, as it were in a cobwebbe, must diligently be searched, and burned from the boughes, before they bring forth other Caterpillars, which doe in December, Ianuary, and February.

Some were wont to break them off, and tread them with their feet, but therewith they be not wholly killed. The fire confumeth all things, and therefore it is

best to burne them.

Against the Pifmires or Ants, when they will hurt the young trees.

Of T the leaves off which are eaten or poisoned of the Ants or Pismires, and where there is any thinge made uncleane in the top of the tree, of those little worms, that rub inpieces with your hands, that it may not sain the other leaves, and that the young sprouts may grow up without any hinderance.

How to keep the Pifmires from the trees.

First make a juice of an herbe, called Portabaca, and mix it with vineger, and sprinkle the stump therwith, or anoint the stumpe with wine dregs. Some take a little weak pitch, but very thin, that it may not hurt the tree.

Another

Another Inftruction.

Take a little bundle of cotton, wooll, flax, or towe, and lay it about the flump, and tie likewife a bundle above, about the flump, and draw it out a little, and the Pifmires can do no hurt: or put about the flump bird-lime.

In what time of the barvest the fruit must be gathered.

The Fruits are not altogether at one timegathered, for they are not ripe all at once, as some pears which shew the ripenesse by the colour, those should be gathered in Summer, and if you let them stand too long, they will not last.

Peares which are ripe in harvest, those may be gathered in O Hober, when the weather is cleare and dry: in harvest in the increase of the Moon, Fruits may be

gathered.

#### **ን**ቀቀቃ ተቀቀቀቀ ተቀቀቀ ተ

Asbort Instruction very profitable and necessary for all those that delight in Gardening to know the times and seasons when it is good to sow and replant all manner of seeds.

Abbages must be sowne in February, March, or April, at the waning of the Moon, and replanted also in the decrease thereof.

Cabbage, Lettuce, in February, March, or Inly, in an old Moon.

Onyons and Leeks must be fowne in February of March, at the waning of the Moon.

D

Bects

The empert Gardener.

Beets must be sown in February, or March, in a

full moone.

Coleworts white and green in February, or March, in an old moone, it is good to replant them.

Parsneps must be sown in February, April or Iune,

alfo in an old moone.

Radish must be sown in February, March, or June, in a new moon.

Pompions must be sown in February, March, or

Innealfo in a new moone.

Cucumbers and Mellons must be sown in Februery, March or June, in an old moone.

Spinage must be fown in February or March, in

an old moone.

Parsley must be sowne in February or March, in a full moone.

Fennell and Anniffeed must be sowne in February

or March, in a full moone.

White Cycory must be sown in February, March, July or August, in a full moon.

Carduus Benedictus must be sowne in February,

March or May, when the moone is old,

Basill must be sowne in March, when the moon is old.

Pursline must be sown in February or March, in a new moone.

Margeram, Violets and Time, must be sowne in February, March or Aprill, in a new moone.

Floure-gentle, Rosemary and Lavender, must be

fowne in February or Aprill, in a new moone.

Rocket and Garden creffes, muß be sowne in Fepruary in a new moon.

Savell

Savell must be sowne in February or March, in a

Saffron must be sown in March, when the moon is old.

Coriander and Borage must be sown in February or March, in a new moon.

Hartshorne and Samphire must be sowne in February, March or Aprill, when the moon is old.

Gilly-flowers, Harts-ease, and Wall-flowers must be sown in March or Aprill, when the moon is old.

Cardons and Artochokes must be sown in Aprill or March when the moon is old.

Chickweed must be sown in February or March, in the full of the moone.

Burnet must be sown in February or March, when the moone is old.

Double Marigolds must be sowne in February or March, in a new moone.

Isopand Savorie must be sowne in March, when the moone is old:

White Poppy must bee sowne in February or March, in a new moone.

Palma Christi must be sowne in February, in a new moone.

Sparages and Sperage is to be fowne in February, when the moone is old.

Larks foot must be sowne in February, when the moone is old.

Note that at all times and feafons, Lettuce, Radish, Spinage and Pafeneps may be fowne.

Note also, from cold are to be kept Coleworts, Cabbage, Lettuce, Bafill, Cardons, Artochokes, and Colefloures.

Da

Worthy

Worthy remedies and focrets availing against the stroying of Snailes, Cankerwormes, the long bodied Moths, Garden-steas, Earthwormes, and Moles.

A fricanus, fingular among the Greek writers of husbandry, reporteth, that Garden plants and roots may well be purged and rid of the harmefull wormes, if their dennes or deep holes be smoaked, the winde aiding, with the dung of the Cow or Oxe

burned.

That worthy Pliny in his first booke of Histories writerh, that if the Owner or Gardener sprinckleth the pure mother of the oyle Olive, without any salt in it, doth also drive the wormes away, and defend the Plants and Herbs from being gnawne of them. And if they shall cleave to the roots of the plants, through malice or breeding of the dung, yet this weedeth them clean away. The plants or herbs will not after be gnawne or harmed by Girden fleas, if with the natural remedy, as with the herbe Rocket, the Gardener shall bestow his beds in many places.

The Coleworts and all pot-herbs are greatly defended from the gnawing of the Garden-fleas, by Radish growing among them. The eager or sharp vineger doth also prevaile, tempered with the juice of Henbane, and sprinkled on the garden fleas. To these, the water in which the herbe Nigella Romana shall be steeped for a night, and sprinkled on the plants, as the Greek Ramphilm reporteth, dothalike

prevaile against the garden fleas.

Paladim Ruthin reporteth, that the notiome vermine or creeping things will not breed of the Pot-

herbs,

herbes, if the Gardener shall before the committing to the earth, dry all the seeds in the skin of the Tortoise, or sow the herbe Mint in many places of the Garden, especially among the Coleworts. The bitter Fitch and Rocket (as I before uttered) bestowed among the pot-herbs, so that the seeds be sowne in the first quarter of the Moon, doe greatly availens. Also the Canker and Palmer wormes, which in many places work great injury both to the Gardens and vines, may the owner or Gardener drive away with the figtree ashes sprinckled on them and the herbs.

There be some which sprinckle the plants and herbes made with the lee of the fig-tree ashes, but it destroyes the wormes, to strew (as experience repor-

teth) the after alone on them.

There be others which rather will to plant or fow that big onyon, named in Latine Scilla or Squilla here and there in beds, or hang them in fundry places of

the Garden.

Others also will to fixe River Cresses with nailes in many places of the garden, which if they shall yet withstand or contend with all these remedies, then may the Gardener apply to exercise this devise, in taking the Ox or Cow urine, and the mother of oyle Olive, which after the well mixing together, and heating over the fire, the same be stirred about untill it be hot, and when through cold, this mixture shall be sprinckled on the pot-herbs and trees, doth marvellously prevaile, as the skilfull anatolius of experience reporteth.

The worthy Paladias Rutilius reporteth, that if the owner or Gardener burne great bundles of the Garlicke blades (without heads) dried, through all the

3 allies

allies of the Garden, and unto these the dung of Backes added, that the savour of the smoke (by the helpe of the wind) may be driven to many places, especially to those where they most abound and swarm, and the Gardener shall see so speedy a de-

ftsuction, as is to be wondred ar.

The worthy Pliny of great knowledge, reporteth that these may be driven from the pot-herbes, if the bitter Fitch feeds be mixed and fown together with them, or the branches of the trees, Crevifes hanged up by the hornes in many places, doth like prevaile. These also are letted from increasing, yea, they in heaps presently gathered are destroied, as the Greeks report of observation, if the Gardener by taking certaine Palmer or Canker-wormes out of the Garden next joyning, shall feeth them in water with Dill. and the same being through cold, shall sprinckle on the herbes and trees, that the mixture may wet and foke through the nefts, even unto the young ones. cleaving together, that they may tafte thereof, will speedily dispatch them. But in this doing, the Gardener must be very wary, and have an attentive eye. that none of the mixture fall on his face or hands.

Besides these, the owner or Gardener may use this remedy certain, and easily prepared, if about the big armes of trees, or stumps of the herbs, he kindle and burn the stronger lime and brimstone together. Or if the owner make a smoake with the Mushromes, growing under the Nut tree, or burne the hoofs of Goats, or the gum Galbanum, or else make a smoake with the Harts horne, the winde aiding, by blowing

towards them.

The husbandmen and gardeners in our turne have found

found out this easie practife, being now commonevery where: which is on this wise, that when these after-showres of rain are cropen into the warm sun, or into places standing against the Sunne, early in the morning shake either their fruits and leaves of the pot-herbes, or the boughes of the trees, for these being yet stiffe, through the cold of the night, are procured of the same, the lighter and sooner to fall, nor able after to recover up againe, so that the Palmer worms thus lying on the ground, are then in a readinesse to be killed of the Gardener.

If the owner mind to destroy any other creeping things noyous to herbes and trees, (which Paladius and Ratilias name, both herb and Leek-wasters) then let him bearken to this invention and devise of the Greek Dyophanes, who willeth to purchale the maw of a Wether sheep new killed, and the same as yet full of his excrementall filth, which lightly cover with the earth in the same place, where these most haunt in the Garden, and after two dayes shall the Gardiner find there, that the mothes with long bodies, and other creeping things will be gathered in divers companies to the place right over it, which the owner shall either remove and carry further, or dig and bury very deep in the same place, that they may not after arise and come forth, which when the Gardener shall have exercised the same but twice or thrice, he shall utterly excinguish, and quite destroy all the kindes of creeping things that anon and spoilthe Garden plants. the wood, which thus

The husbandmen in Flanders arme the stockes, and compasse the bigger armes of their trees, with wisps of straw handsomely made and fastened or bound.

bound about by which the Palmer wormer are confrained to creep up to the tops of the trees, and there staid, fo that, (as it were by fnares and engines laid) thefe in the end are driven away, or thus in their way begun, are speedily by soone after procured to turne backe againe; As anto the remedies of the Snalles particularly belongs. These may the Gardener likewife chafe from the kirchin herbs, if he either fprinckle the new mother of the cyle olive, or foot of the chimney on the herbs, as if he bestowed the bitter fitch in beds among them, which also availes again & other noisome worms, and creeping things, as I afore ucce red, that if the Gardener would possesse a greene and delectable Garden, let him then sprinckle diligently all the quarters, beds, and borders of the Garden, with the mixture of water and powder of Fennis greeke tempered rogether, or fer upright in the midde of the Garden, the whole bare head without the flesh of the unchaste Asse, as I afore wrote.

### Excellent inventions and helps against the Garden Moles.

The skilfull Paxanum hath left in writing, that if the Gardener should make hollow a big nut, or bore a hollow hole into some sound piece of wood being narrow, in filling the one or the other with Rosin, Pitch, Chasse and Brimstone, of each so much as shall suffice to the filling of the Nut, or hollow hole in the wood, which thus prepared in a readinesse, stop every where with diligence, all the goings forth, and breathing holes of the Mole, that by those the suming smoke in no manner may liste our, yet

fo handle the matter, that one mouth and hole bee only left open, and the fame so large, that well the nut or vessell kindled within, may be laid within the mouth of it, whereby it may take the wind of the one side, which may so send in the savour both of the rossin and brimstone into the hollow tombe, or resting place of the Mole; by the same practise so worker, manly handled, by filling the holes with the smoak, shall the owner or Gardener either drive quite away all the Moles in the ground, or finde them in a short time dead.

There be some that take the white Neesewort, or the rinde of Cynecrambes beaten and farced, and with Barley meale and egs finely tempered together, they make both Cakes and Pasties wrought with wine and milk, and those they lay within the Moles den or hole.

Albert w of worthy memory reporteth, that if the owner or Gardener closeth or diligently stoppeth the mouths of the Moleholes, with the garlicke, onyon, or leek, it shall either drive the Moles away, or kil them through the strong favour stinking or brea-

thing into them.

Many there be, that to drive away these harmfull Moles, doe bring up young Cats in their Garden ground, and make tame Weasels, to the end, that either of these through the hunting of them, may so drive away this pestiferous anoyance, being taught to watch at their strair passages, and mouths of the holes comming forth.

Others there be also, which differely fill and stop up their holes with the red Okare or Ruddell, and juice of the wild Cucumber, or sow the seeds of The expert Guranner.

Patho Chrift, being a kind of Sayrion, in beds, through which they will not after cast up, nor earry thereabout.

But some exercise this easie practise, in taking a live Mole, and burning the pouder of brimstone abour him, being in a deep earthen por, through which he is procured to cry, all others in the meane time as they report, are moved to resort thither.

There are some besides, which lay silke snares at

the mouth of their holes.


To the simple Husbandmen may this easie pradife of no cost suffice, in setting downe into the earth, a stiffe rod or green branch of the Eldertree.


# FINIS

erd can sythis pellistrens aroganes, being tangue to watch of their fleair passess, and mouths of the holes contain a lottle.


Lop up their boles with the red Okure or Ruddell, and juice of the wild Cucumber, or for the feeds of Palms.

The espery Gardener.






A Direction to fet or lay your lines or thread to make or draw a simple Knot, without a border.




You must leave your Lines as they be first ser, untill your Knct be altogether finished or done.

The Manner or Ordering to fet the thread or



A Direction to faften your Lines to make another manner of Knot.




UM

The Tempers Gardener

31


A plaine Knot without Lines.



A plaine Knot without Lines A





# Another plaine Knot withour Lines.




The supert Gardener?


#### Anochen.







41






45







A Direction to fasten Cords or Lines to draw a
Knot with a Border; as also to make a Border
of Beds parted in the midst.




A Direction of the Cords falfned upon the Border, with a Knot in the midft.




# A Border with a Knot in the midst thereof.




A Border or Knot divided or parted, containing five small Knots.



The forme of the Lines fet upon the Knot, whose Squares or Beds are parted.




A Border of Beds or Squares parted; and the midst thereof.



# The imper Gurdener T





.

60)